

Reducing the Number of Homogeneous Linear Equations in Finding Annihilators

Deepak Kumar Dalai and Subhamoy Maitra

Applied Statistics Unit, Indian Statistical Institute,
203, B T Road, Calcutta 700 108, INDIA
{deepak_r, subho}@isical.ac.in

Abstract. Given a Boolean function f on n -variables, we find a reduced set of homogeneous linear equations by solving which one can decide whether there exist annihilators at degree d or not. Using our method the size of the associated matrix becomes $\nu_f \times (\sum_{i=0}^d \binom{n}{i} - \mu_f)$, where, $\nu_f = |\{x | wt(x) > d, f(x) = 1\}|$ and $\mu_f = |\{x | wt(x) \leq d, f(x) = 1\}|$ and the time required to construct the matrix is same as the size of the matrix. This is a preprocessing step before the exact solution strategy (to decide on the existence of the annihilators) that requires to solve the set of homogeneous linear equations (basically to calculate the rank) and this can be improved when the number of variables and the number of equations are minimized. As the linear transformation on the input variables of the Boolean function keeps the degree of the annihilators invariant, our preprocessing step can be more efficiently applied if one can find an affine transformation over $f(x)$ to get $h(x) = f(Bx + b)$ such that $\mu_h = |\{x | h(x) = 1, wt(x) \leq d\}|$ is maximized (and in turn ν_h is minimized too). We present an efficient heuristic towards this. Our study also shows for what kind of Boolean functions the asymptotic reduction in the size of the matrix is possible and when the reduction is not asymptotic but constant.

Keywords: Algebraic Attacks, Algebraic Normal Form, Annihilators, Boolean Functions, Homogeneous Linear Equations.

1 Introduction

Results on algebraic attacks have received a lot of attention recently in studying the security of crypto systems [2, 4, 6, 9, 11–15, 22, 1, 21, 16]. Boolean functions are important primitives to be used in the crypto systems and in view of the algebraic attacks, the annihilators of a Boolean function play considerably serious role [5, 7, 10, 17–19, 23, 24].

Denote the set of all n -variable Boolean functions by B_n . One may refer to [17] for the detailed definitions related to Boolean functions, e.g., truth table, algebraic normal form (ANF), algebraic degree (deg), weight (wt), nonlinearity (nl) and Walsh spectrum of a Boolean function. Any Boolean function can

be uniquely represented as a multivariate polynomial over $GF(2)$, called the algebraic normal form (ANF), as

$$f(x_1, \dots, x_n) = a_0 + \sum_{1 \leq i \leq n} a_i x_i + \sum_{1 \leq i < j \leq n} a_{i,j} x_i x_j + \dots + a_{1,2,\dots,n} x_1 x_2 \dots x_n,$$

where the coefficients $a_0, a_i, a_{i,j}, \dots, a_{1,2,\dots,n} \in \{0, 1\}$. The algebraic degree, $\deg(f)$, is the number of variables in the highest order term with non zero coefficient. Given $f \in B_n$, a nonzero function $g \in B_n$ is called an annihilator of f if $f * g = 0$. A function f should not be used if f or $1 + f$ has a low degree annihilator. It is also known [14, 23] that for any function f or $1 + f$ must have an annihilator at the degree $\lceil \frac{n}{2} \rceil$. Thus the target of a good design is to use a function f such that neither f nor $1 + f$ has an annihilator at a degree less than $\lceil \frac{n}{2} \rceil$. Thus there is a need to construct such functions and the first one in this direction appeared in [18]. Later symmetric functions with this property has been presented in [19] followed by [7]. However, all these constructions are not good in terms of other cryptographic properties.

Thus there is a need to study the Boolean functions, which are rich in terms of other cryptographic properties, in terms of their annihilators. One has to find out the annihilators of a given Boolean function for this. Initially a basic algorithm in finding the annihilators has been proposed in [23, Algorithm 2]. A minor modification of [23, Algorithm 2] has been presented very recently in [8] to find out relationships for algebraic and fast algebraic attacks. In [7], there is an efficient algorithm to find the annihilators of symmetric Boolean functions, but symmetric Boolean functions are not cryptographically promising. Algorithms using Gröbner bases are also interesting in this area [3], but still they are not considerably efficient. Recently more efficient algorithms have been designed in this direction [1, 21]. The algorithm presented in [1] can be used efficiently to find out relationships for algebraic and fast algebraic attacks. In [1], matrix triangularization has been exploited nicely to solve the annihilator finding problem (of degree d for an n -variable function) in $O(\binom{n}{d}^2)$ time complexity. In [21] a probabilistic algorithm having time complexity $O(n^d)$ has been proposed where the function is divided to its sub functions recursively and the annihilators of the sub functions are checked to study the annihilators of the original function.

The main idea in our effort is to reduce the size of the matrix (used to solve the system of homogeneous linear equations) as far as possible, which has not yet been studied in a disciplined manner to the best of our knowledge. We could successfully improve the handling of equations associated with small weight inputs of the Boolean function. This uses certain structure of the matrix that we discover here. We start with a matrix $M_{n,d}(g)$ (see Theorem 1) which is self inverse and its discovered structure allows to compute the new equations efficiently by considering the matrix UA^r (see Theorem 3 in Section 3). Moreover, each equation associated with a low weight input point directly provides the value of an unknown coefficient of the annihilator, which is the key point that allows to lower the number of unknowns. Further reduction in the size of the matrix

is dependent on getting a proper linear transformation on the input variables of the Boolean function, which is discussed in Section 4.

One may wonder whether the very recently available strategies in [1, 21] can be applied after the initial reduction proposed in this paper to get further improvements in finding the lowest degree annihilators. The standard Gaussian reduction technique ([21, Algorithm 1]) is used in the main algorithm [21, Algorithm 2], and in that case our idea of reduction of the matrix size will surely provide improvement. However, the ideas presented in [1, Algorithm 1, 2] and [21, Algorithm 3] already exploit the structure of the linear system in an efficient way. In particular, the algorithms in [1] by themselves deal with the equations of small weight efficiently. Thus it is not clear whether the reduction of matrix size proposed by us can be applied to exploit further efficiency from these algorithms.

2 Preliminaries

Consider all the n -variable Boolean functions of degree at most d , i.e., $\mathcal{R}(n, d)$, the Reed-Muller code of order d and length 2^n . Any Boolean function can be seen as a multivariate polynomial over $GF(2)$. Note that $\mathcal{R}(n, d)$ is a vector subspace of the vector space \mathcal{B}_n , the set of all n -variable Boolean functions. Now if we consider the elements of $\mathcal{R}(n, d)$ as the multivariate polynomials over $GF(2)$, then the standard basis is the set of all nonzero monomials of degree $\leq d$. That is, the standard basis is

$$S_{n,d} = \{x_{i_1} \dots x_{i_l} : 1 \leq l \leq d \text{ and } 1 \leq i_1 < i_2 < \dots < i_l \leq n\} \cup \{1\},$$

where the input variables of the Boolean functions are x_1, \dots, x_n .

The ordering among the monomials is considered in lexicographic ordering ($<_l$) as usual, i.e., $x_{i_1}x_{i_2} \dots x_{i_k} <_l x_{j_1}x_{j_2} \dots x_{j_l}$ if either $k < l$ or $k = l$ and there is $1 \leq p \leq k$ such that $i_k = j_k$, $i_{k-1} = j_{k-1}, \dots, i_{p+1} = j_{p+1}$ and $i_p < j_p$. So, the set $S_{n,d}$ is a totally ordered set with respect to this lexicographical ordering ($<_l$). Using this ordering we refer the monomials according their order, i.e., the k -th monomial as m_k , $1 \leq k \leq \sum_{i=0}^d \binom{n}{i}$ following the convention $m_l <_l m_k$ if $l < k$.

Definition 1. Given $n > 0$, $0 \leq d \leq n$, we define a mapping

$$v_{n,d} : \{0, 1\}^n \mapsto \{0, 1\}^{\sum_{i=0}^d \binom{n}{i}},$$

such that $v_{n,d}(x) = (m_1(x), m_2(x), \dots, m_{\sum_{i=0}^d \binom{n}{i}}(x))$. Here $m_i(x)$ is the i th monomial as in the lexicographical ordering ($<_l$) evaluated at the point $x = (x_1, x_2, \dots, x_n)$.

To evaluate the value of the t -th coordinate of $v_{n,d}(x_1, x_2, \dots, x_n)$ for $1 \leq t \leq \sum_{i=0}^d \binom{n}{i}$, i.e., $[v_{n,d}(x_1, \dots, x_n)]_t$, one requires to calculate the value of the monomial m_t (either 0 or 1) at (x_1, x_2, \dots, x_n) . Now we define a matrix $M_{n,d}$ with respect to a n -variable function f . To define this we need another similar

ordering ($<^l$) over the elements of vector space $\{0, 1\}^n$. We say for $u, v \in \{0, 1\}^n$, $u <^l v$ if either $wt(u) < wt(v)$ or $wt(u) = wt(v)$ and there is some $1 \leq p \leq n$ such that $u_n = v_n, u_{n-1} = v_{n-1}, \dots, u_{p+1} = v_{p+1}$ and $u_p = 0, v_p = 1$.

Definition 2. Given $n > 0$, $0 \leq d \leq n$ and an n -variable Boolean function f , we define a $wt(f) \times \sum_{i=0}^d \binom{n}{i}$ matrix

$$M_{n,d}(f) = \begin{bmatrix} v_{n,d}(X_1) \\ v_{n,d}(X_2) \\ \vdots \\ v_{n,d}(X_{wt(f)}) \end{bmatrix}$$

where any X_i is an n -bit vector and $supp(f) = \{X_1, X_2, \dots, X_{wt(f)}\}$ and $X_1 <^l X_2 <^l \dots <^l X_{wt(f)}$; $supp(f)$ is the set of inputs for which f outputs 1.

Note that the matrix $M_{n,d}(f)$ is the transpose of the restricted generator matrix for Reed-Muller code of length 2^n and order d , $\mathcal{R}(d, n)$, to the support of f (see also [9, Page 7]). Any row of the matrix $M_{n,d}(f)$ corresponding to an input vector (x_1, \dots, x_n) is

$$\underbrace{1}_{0 \text{ deg}} \underbrace{x_1, \dots, x_i, \dots, x_n}_{1 \text{ deg}} \dots \underbrace{x_1 \dots x_d, \dots, x_{i_1} \dots x_{i_d}, \dots, x_{n-d+1} \dots x_n}_{d \text{ deg}}.$$

Each column of the matrix is represented by a specific monomial and each entry of the column tells whether that monomial is satisfied by the input vector which identifies the row, i.e., the rows of this matrix correspond to the evaluations of the monomials having degree at most d on support of f . As already discussed, here we have one-to-one correspondence from the input vectors $x = (x_1, \dots, x_n)$ to the row vectors $v_{n,d}(x)$ of length $\sum_{i=0}^d \binom{n}{i}$. So, each row is fixed by an input vector.

2.1 Annihilator of f and rank of the matrix $M_{n,d}(f)$

Let f be an n -variable Boolean function. We are interested to find out the lowest degree annihilators of f . Let $g \in B_n$ be an annihilator of f , i.e., $f(x_1, \dots, x_n) * g(x_1, \dots, x_n) = 0$. In terms of truth table, this means that the function f AND g will be a constant zero function, i.e., for each vector $(x_1, \dots, x_n) \in \{0, 1\}^n$, the output of f AND g will be zero. That means,

$$g(x_1, \dots, x_n) = 0 \text{ if } f(x_1, \dots, x_n) = 1. \quad (1)$$

Suppose degree of the function g is $\leq d$, then the ANF of g is of the form $g(x_1, \dots, x_n) = a_0 + \sum_{i=0}^n a_i x_i + \dots + \sum_{1 \leq i_1 < i_2 < \dots < i_d \leq n} a_{i_1, \dots, i_d} x_{i_1} \dots x_{i_d}$ where the subscripted a 's are from $\{0, 1\}$ and not all of them are zero. Following Equation 1, we get the following $wt(f)$ many homogeneous linear equations

$$a_0 + \sum_{i=0}^n a_i x_i + \dots + \sum_{1 \leq i_1 < i_2 < \dots < i_d \leq n} a_{i_1, \dots, i_d} x_{i_1} \dots x_{i_d} = 0, \quad (2)$$

considering the input vectors $(x_1, \dots, x_n) \in \text{supp}(f)$. This is a system of homogeneous linear equations on a 's with $\sum_{i=0}^d \binom{n}{i}$ many a 's as variables. The matrix form of this system of equations is $M_{n,d}(f) A^{tr} = O$, where $A = (a_0, a_1, a_2, \dots, a_{n-d+1}, \dots, a_n)$, the row vector of coefficients of the monomials which are ordered according to the lexicographical order $<_l$. Each nonzero solution of the system of equations formed by Equation 2 gives an annihilator g of degree $\leq d$. This is basically the Algorithm 1 presented in [23]. Since the number of solutions of this system of equations are connected to the rank of the matrix $M_{n,d}(f)$, it is worth to study the rank and the set of linear independent rows/columns of matrix $M_{n,d}(f)$. If the rank of matrix $M_{n,d}(f)$ is equal to $\sum_{i=0}^d \binom{n}{i}$ (i.e., number of columns) then the only solution is the zero solution. So, for this case f has no annihilator of degree $\leq d$. This implies that the number of rows \geq number of columns, i.e., $wt(f) \geq \sum_{i=0}^d \binom{n}{i}$ which is the Theorem 1 in [17]. If the rank of matrix is equal to $\sum_{i=0}^d \binom{n}{i} - k$ for $k > 0$ then the number of linearly independent solutions of the system of equations is k which gives k many linearly independent annihilators of degree $\leq d$ and $2^k - 1$ many number of annihilators of degree $\leq d$. However, to implement algebraic attack one needs only linearly independent annihilators. Hence, finding the degree of lowest degree annihilator of either f or $1 + f$, one can use the following algorithm.

Algorithm 1

```

for( $i = 1$  to  $\lceil \frac{n}{2} \rceil - 1$ ) {
    find the rank  $r_1$  of the matrix  $M_{n,i}(f)$ ;
    find the rank  $r_2$  of the matrix  $M_{n,i}(1 + f)$ ;
    if  $\min\{r_1, r_2\} < \sum_{j=0}^i \binom{n}{j}$  then output  $i$ ;
}
output  $\lceil \frac{n}{2} \rceil$ ;

```

Since either f or $1 + f$ has an annihilator of degree $\leq \lceil \frac{n}{2} \rceil$, we are interested only to check till $i = \lceil \frac{n}{2} \rceil$. This algorithm is equivalent to Algorithm 1 in [23].

The simplest and immediate way to solve the system of these equations or find out the rank of $M_{n,d}(f)$, $M_{n,d}(1 + f)$ is the Gaussian elimination process. To check the existence or to enumerate the annihilators of degree $\leq \lceil \frac{n}{2} \rceil$ for a balanced function, the complexity is approximately $(2^{n-2})^3$. Considering this time complexity, it is not encouraging to check annihilators of a function of 20 variables or more using the presently available computing power. However, given n and d , the matrix $M_{n,d}(f)$ has pretty good structure, which we explore in this paper towards a better algorithm (that is solving the set of homogeneous linear equations in an efficient way by decreasing the size of the matrix involved).

3 Faster strategy to construct the set of homogeneous linear equations

In this section we present an efficient strategy to reduce the set of homogeneous linear equations. First we present a technical result.

Theorem 1. Let g be an n -variable Boolean function defined as $g(x) = 1$ iff $wt(x) \leq d$ for $0 \leq d \leq n$. Then $M_{n,d}(g)^{-1} = M_{n,d}(g)$, i.e., $M_{n,d}(g)$ is a self inverse matrix.

Proof. Suppose $\mathcal{F} = M_{n,d}(g)M_{n,d}(g)$. Then the i -th row and j -th column entry of \mathcal{F} (denoted by $\mathcal{F}_{i,j}$) is the scalar product of i -th row and j -th column of $M_{n,d}(g)$. Suppose the i -th row is $v_{n,d}(x)$ for $x \in \{0,1\}^n$ having x_{q_1}, \dots, x_{q_l} as 1 and others are 0. Further consider that the j -th column is the evaluation of the monomial $x_{r_1} \dots x_{r_k}$ at the input vectors belonging to the support of g . If $\{r_1, \dots, r_k\} \not\subseteq \{q_1, \dots, q_l\}$ then $\mathcal{F}_{i,j} = 0$. Otherwise, $\mathcal{F}_{i,j} = \binom{l-k}{0} + \binom{l-k}{1} + \dots + \binom{l-k}{l-k} \mod 2 = 2^{l-k} \mod 2$. So, $\mathcal{F}_{i,j} = 1$ iff $\{x_{r_1}, \dots, x_{r_k}\} = \{x_{q_1}, \dots, x_{q_l}\}$. That implies, $\mathcal{F}_{i,j} = 1$ iff $i = j$ i.e., \mathcal{F} is identity matrix. Hence, $M_{n,d}(g)$ is its own inverse. \square

See the following example for the structure of $M_{n,d}(g)$ when $n = 4$ and $d = 2$.

Example 1. Let us present an example of $M_{n,d}(g)$ for $n = 4$ and $d = 2$. We have $\{1, x_1, x_2, x_3, x_4, x_1x_2, x_1x_3, x_2x_3, x_1x_4, x_2x_4, x_3x_4\}$, the list of 4-variable monomials of degree ≤ 2 in ascending order ($<_l$).

Similarly, $\{(0,0,0,0), (1,0,0,0), (0,1,0,0), (0,0,1,0), (0,0,0,1), (1,1,0,0), (1,0,1,0), (0,1,1,0), (1,0,0,1), (0,1,0,1), (0,0,1,1)\}$ present the 4 dimensional vectors of weight ≤ 2 in ascending order ($<^l$). So the matrix

$$M_{4,2}(g) = \begin{bmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 1 & 1 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 1 & 0 & 1 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 1 & 1 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 & 1 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 1 & 0 & 1 & 0 & 1 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 1 & 1 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \end{bmatrix}$$

One may check that $M_{4,2}(g)$ is self inverse.

Lemma 1. Let A be a nonsingular $m \times m$ binary matrix where the row vectors are denoted as v_1, v_2, \dots, v_m . Let U be a $k \times m$ binary matrix, $k \leq m$, where the rows are denoted as u_1, u_2, \dots, u_k . Let $W = UA^{-1}$, a $k \times m$ binary matrix. Consider that a matrix A' is formed from A by replacing the rows $v_{i_1}, v_{i_2}, \dots, v_{i_k}$ of A by the vectors u_1, u_2, \dots, u_k . Further consider that a $k \times k$ matrix W' is formed by taking the i_1 -th, i_2 -th, \dots, i_k -th columns of W (out of m columns). Then A' is nonsingular iff W' is nonsingular.

Proof. Without loss of generality, we can take $i_1 = 1, i_2 = 2, \dots, i_k = k$. So, the row vectors of A' are $u_1, \dots, u_k, v_{k+1}, \dots, v_m$.

We first prove that if the row vectors of A' are not linearly independent then the row vectors of W' are also not linearly independent. As the row vectors of A' are not linearly independent, we have $\alpha_1, \alpha_2, \dots, \alpha_m \in \{0,1\}$ (not all zero) such that $\sum_{i=1}^k \alpha_i u_i + \sum_{i=k+1}^m \alpha_i v_i = 0$. If $\alpha_i = 0$ for all i , $1 \leq i \leq k$ then $\sum_{i=k+1}^m \alpha_i v_i = 0$ which implies $\alpha_i = 0$ for all i , $k+1 \leq i \leq m$ as $v_{k+1}, v_{k+2}, \dots, v_m$ are linearly independent. So, all α_i 's for $1 \leq i \leq k$ can not be zero.

Further, we have $UA^{-1} = W$, i.e., $U = WA$, i.e.,

$$\begin{pmatrix} u_1 \\ u_2 \\ \vdots \\ u_k \end{pmatrix} = \begin{pmatrix} w_1 \\ w_2 \\ \vdots \\ w_k \end{pmatrix} \begin{pmatrix} v_1 \\ v_2 \\ \vdots \\ v_m \end{pmatrix}, \text{ i.e., } u_i = w_i \begin{pmatrix} v_1 \\ v_2 \\ \vdots \\ v_m \end{pmatrix}.$$

$$\text{Hence, } \sum_{i=1}^k \alpha_i u_i = \sum_{i=1}^k \alpha_i w_i \begin{pmatrix} v_1 \\ v_2 \\ \vdots \\ v_m \end{pmatrix} = r \begin{pmatrix} v_1 \\ v_2 \\ \vdots \\ v_m \end{pmatrix}$$

where $r = (r_1, r_2, \dots, r_m) = \sum_{i=1}^k \alpha_i w_i$.

If the restricted matrix W' were nonsingular, the vector $r' = (r_1, r_2, \dots, r_k)$ is non zero as $(\alpha_1, \alpha_2, \dots, \alpha_k)$ is not all zero. Hence, $\sum_{i=1}^k \alpha_i u_i + \sum_{i=k+1}^m \alpha_i v_i = 0$, i.e., $\sum_{i=1}^k r_i v_i + \sum_{i=k+1}^m (r_i + \alpha_i) v_i = 0$. This contradicts that v_1, v_2, \dots, v_m are linearly independent as $r' = (r_1, r_2, \dots, r_k)$ is nonzero. Hence W' must be singular. This proves one direction.

On the other direction if the restricted matrix W' is singular then there are $\beta_1, \beta_2, \dots, \beta_k$ not all zero such that $\sum_{i=0}^k \beta_i w_i = (0, \dots, 0, s_{k+1}, \dots, s_m)$. Hence,

$$\sum_{i=0}^k \beta_i u_i = \sum_{i=1}^k \beta_i w_i \begin{pmatrix} v_1 \\ v_2 \\ \vdots \\ v_m \end{pmatrix} = s_{k+1} v_{k+1} + \dots + s_m v_m, \text{ i.e., } \sum_{i=0}^k \beta_i u_i +$$

$\sum_{i=k+1}^m s_i v_i = 0$ which says matrix A' is singular. \square

Following Lemma 1, one can check the nonsingularity of the larger matrix A' by checking the nonsingularity of the reduced matrix W' . Thus checking the nonsingularity of the larger matrix A' will be more efficient if the computation of matrix product $W = UA^{-1}$ can be done efficiently. The self inverse nature of the matrix $M_{n,d}(g)$ presented in Theorem 1 helps to achieve this efficiency. In the rest of this section we will study this in detail. In the following result we present the Lemma 1 in more general form.

Theorem 2. *Let A be a nonsingular $m \times m$ binary matrix with m -dimensional row vectors v_1, v_2, \dots, v_m and U be a $k \times m$ binary matrix with m -dimensional row vectors u_1, u_2, \dots, u_k . Consider $W = UA^{-1}$, a $k \times m$ matrix. The matrix A' , formed from A by removing the rows $v_{i_1}, v_{i_2}, \dots, v_{i_l}$ ($l \leq m$) from A and adding the rows u_1, u_2, \dots, u_k ($k \geq l$), is of rank m iff the rank of restricted $k \times l$ matrix W' including only the i_1 -th, i_2 -th, \dots , i_l -th columns of W is l .*

Proof. Here, the rank of matrix W' is l . So, there are l many rows of W' , say $w'_{p_1}, \dots, w'_{p_l}$ which are linearly independent. So, following the Lemma 1 we have the matrix A'' formed by replacing the rows v_{i_1}, \dots, v_{i_l} of A by u_{p_1}, \dots, u_{p_l} is nonsingular, i.e., rank is m . Hence the matrix A' where some more rows are added to A'' has rank m . The other direction can also be shown similar to the proof of the other direction in Lemma 1. \square

Now using Theorem 1 and Theorem 2, we describe a faster algorithm to check the existence of annihilators of certain degree d of a Boolean function f . Suppose g be the Boolean function described in Theorem 1, i.e., $\text{supp}(g) = \{x | 0 \leq \text{wt}(x) \leq d\}$. In Theorem 1, we have already shown that $M_{n,d}(g)$ is nonsingular matrix (in fact it is self inverse). Let $\{x | \text{wt}(x) \leq d \text{ and } f(x) = 0\} = \{x_1, x_2, \dots, x_l\}$ and $\{x | \text{wt}(x) > d \text{ and } f(x) = 1\} = \{y_1, y_2, \dots, y_k\}$. Then we consider $M_{n,d}(f)$ as A , $v_{n,d}(x_1), \dots, v_{n,d}(x_l)$ as v_{i_1}, \dots, v_{i_l} and $v_{n,d}(y_1), \dots, v_{n,d}(y_k)$ as u_1, \dots, u_k . Then following Theorem 2 we can ensure whether $M_{n,d}(f)$ is nonsingular. If it is nonsingular, then there is no annihilator of degree $\leq d$, else there are annihilator(s). We may write this in a more concrete form as the following corollary to Theorem 2.

Corollary 1. *Let f be an n -variable Boolean function. Let A^r be the restricted matrix of $A = M_{n,d}(g)$, by taking the columns corresponding to the monomials $x_{i_1}x_{i_2}\dots x_{i_l}$ such that $l \leq d$ and $f(x) = 0$ when $x_{i_1} = 1, x_{i_2} = 1, \dots, x_{i_l} = 1$ and*

rest of the input variables are 0. Further $U = \begin{pmatrix} v_{n,d}(y_1) \\ v_{n,d}(y_2) \\ \vdots \\ v_{n,d}(y_k) \end{pmatrix}$, where $\{y_1, \dots, y_k\} = \{x | \text{wt}(x) > d \text{ and } f(x) = 1\}$. If rank of UA^r is l then there is no annihilator of degree $\leq d$, else there are annihilator(s) of degree $\leq d$.

Proof. As per Theorem 2, here $W = UA^{-1} = UA$, since A is self inverse, following Theorem 1 and hence W' is basically UA^r . Thus the proof follows. \square

Now we can use the following technique for fast computation of the matrix multiplication UA^r . For this we first present a technical result and its proof is similar in the line of the proof of Theorem 1.

Proposition 1. *Consider g as in Theorem 1. Let $y \in \{0,1\}^n$ such that i_1, i_2, \dots, i_p -th places are 1 and other places are 0. Consider the j -th monomial $m_j = x_{j_1}x_{j_2}\dots x_{j_q}$ according the ordering $<_l$. Then the j -th entry of $v_{n,d}(y)M_{n,d}(g)$ is 0 if $\{j_1, \dots, j_q\} \not\subseteq \{i_1, \dots, i_p\}$ else the value is $\sum_{i=0}^{d-q} \binom{p-q}{i} \bmod 2$.*

Following Proposition 1, we can get each row of U as some $v_{n,d}(y)$ and each column of A^r as m_j and construct the matrix UA^r . One can precompute the sums $\sum_{i=0}^{d-q} \binom{p-q}{i} \bmod 2$ for $d+1 \leq p \leq n$ and $0 \leq q \leq d$, and store them and the total complexity for calculating them is $O(d^2(n-d))$. These sums will be used to fill up the matrix UA^r which is an $l \times k$ matrix according to Corollary 1. Let us denote $\mu_f = |\{x | \text{wt}(x) \leq d, f(x) = 1\}|$ and $\nu_f = |\{x | \text{wt}(x) > d, f(x) = 1\}|$. Then $\text{wt}(f) = \mu_f + \nu_f$ and the matrix UA^r is of dimension $\nu_f \times (\sum_{i=0}^d \binom{n}{i} - \mu_f)$. Clearly $O(d^2(n-d))$ can be neglected with respect to $\nu_f \times (\sum_{i=0}^d \binom{n}{i} - \mu_f)$. Thus we have the following result.

Theorem 3. *Consider U and A^r as in Corollary 1. The time (and also space) complexity to construct the matrix UA^r is of the order of $\nu_f \times (\sum_{i=0}^d \binom{n}{i} - \mu_f)$. Further checking the rank of UA^r (as given in Corollary 1) one can decide whether f has an annihilator at degree d or not.*

In fact, to check the rank of the matrix UA^r using Gaussian elimination process, we need not store the ν_f many rows at the same time. One can add one row (following the calculation to compute a row of the matrix given in Proposition 1) at a time incrementally to the previously stored linearly independent rows by checking whether the present row is linearly independent with respect to the already stored rows. If the current row is linearly independent with the existing ones, then we do row operations and add the new row to the previously stored matrix. Otherwise we reject the new row. Hence, our matrix size never crosses the size $(\sum_{i=0}^d \binom{n}{i} - \mu_f) \times (\sum_{i=0}^d \binom{n}{i} - \mu_f)$.

If ν_f (the number of rows) is less than $(\sum_{i=0}^d \binom{n}{i} - \mu_f)$ (the number of variables), then there will be nontrivial solutions and we can directly say that the annihilators exist. Thus we always need to concentrate on the case $\nu_f \geq (\sum_{i=0}^d \binom{n}{i} - \mu_f)$, where the matrix size $(\sum_{i=0}^d \binom{n}{i} - \mu_f) \times (\sum_{i=0}^d \binom{n}{i} - \mu_f)$ provides a further reduction than the matrix size $\nu_f \times (\sum_{i=0}^d \binom{n}{i} - \mu_f)$ and one can save more space. This will be very helpful when one tries to check the annihilators of small degree d .

Refer to Subsection 3.1 below for detailed description that this algorithm provides asymptotic improvement than [23] in terms of constructing this reduced set of homogeneous linear equations. In terms of the overall algorithm to find the annihilators, our algorithm works around eight times further than [23] in general. Using our strategy to find the reduced matrix first and then using the standard Gaussian elimination technique, we could find the annihilators of any random balanced Boolean functions on 16 variables in around 2 hours in a Pentium 4 personal computer with 1 GB RAM. Note that, the very recently known efficient algorithms [1, 21] can work till 20 variables.

3.1 Comparison with Meier et al [23] algorithm

Here we compare the time and space complexity of our strategy with [23, Algorithm 2]. In paper [23], Algorithm 2 is probabilistic. In this draft we study the time and space complexity of the algorithm along with its deterministic version. Using these algorithms we check whether there exist annihilators of degree $\leq d$ of an n -variable function f . As we have already described, ANF of any n -variable function g of degree d is of the form $g(x_1, \dots, x_n) = a_0 + \sum_{i=0}^n a_i x_i + \dots + \sum_{1 \leq i_1 < i_2 < \dots < i_d \leq n} a_{i_1, \dots, i_d} x_{i_1} \dots x_{i_d}$. First we present the exact probabilistic algorithm [23, Algorithm 2].

Algorithm 2

1. Initialize weight $w = 0$.
2. For all x 's of weight w with $f(x) = 1$, substitute each x in $g(x) = 0$ to derive a linear equation on the coefficients of g , with a single coefficient of weight w . Use this equation to express this coefficient iteratively by coefficients of lower weight.
3. If $w < d$, increment w by 1 and go to step 2.

4. Choose random arguments x of arbitrary weight such that $f(x) = 1$ and substitute in $g(x) = 0$, until there are same number of equations as unknowns.
5. Solve the linear system. If there is no solution, output no annihilator of degree d , but if there is a solution then it is not clear whether there is an annihilator of degree d or not.

Next we present the deterministic version of the original probabilistic algorithm [23, Algorithm 2].

Algorithm 3

1. Initialize weight $w = 0$.
2. For all x 's of weight w with $f(x) = 1$, substitute each x in $g(x) = 0$ to derive a linear equation in the coefficients of g , with a single coefficient of weight w . Use this equation to express this coefficient iteratively by coefficients of lower weight.
3. If $w < d$, increment w by 1 and go to step 2.
4. Substitute x such that $wt(x) > d$ and $f(x) = 1$ in $g(x) = 0$ to get linear equation in the coefficient of g .
5. Solve the linear system. Output no annihilator of degree d iff there is no non zero solution.

Since first three steps of both algorithms are same, we initially study the time and space complexity of both the algorithms for first three steps for a randomly choosen balanced function f . In step 2, we apply x , such that weight of $x \leq d$ and $f(x) = 1$, in $g(x)$ and hence we get a linear equation in the coefficient of g such that a single coefficient of that weight is expressed as linear combination of its lower weight coefficients. Here we consider a particular w for each iteration. As f is random and balanced, one can expect that there are $\frac{1}{2}\binom{n}{w}$ many input vectors of weight w in set $\text{supp}(f)$. For each $x = (x_1, \dots, x_n) \in \text{supp}(f)$ where x_{i_1}, \dots, x_{i_w} are 1 and others are 0 of weight w , we will get linear equation of the form

$$a_{i_1, \dots, i_w} = a_0 + \sum_{j=1}^w a_{i_j} + \dots + \sum_{\{k_1, \dots, k_{w-1}\} \subset \{i_1, \dots, i_w\}} a_{k_1, \dots, k_{w-1}}. \quad (3)$$

To store one equation we need $\sum_{i=0}^w \binom{n}{i}$ many memory bits (some places will be 0, some will be 1). There are $\sum_{i=0}^{w-1} \binom{w}{i}$ many coefficients in the right hand side of the Equation 3. As f is random, one can expect that half of them can be eliminated using the equations obtained by lower weight input support vectors. So, $\sum_{i=0}^w \binom{n}{i} + \frac{1}{2} \sum_{i=0}^{w-1} \left(\binom{w}{i} \sum_{j=0}^{i-1} \binom{n}{j} \right)$ order of computation is required to establish an equation. Here w varies from 0 to d and there are approximately $\frac{1}{2} \sum_{w=0}^d \binom{n}{w}$ many support vectors of weight $\leq d$. Hence at the starting of step 4 the space complexity is $S1 = \frac{1}{2} \sum_{w=0}^d \left(\binom{n}{w} \sum_{i=0}^w \binom{n}{i} \right)$ and time complexity is $T1 = \frac{1}{2} \sum_{w=0}^d \left(\binom{n}{w} \left(\sum_{i=0}^w \binom{n}{i} + \frac{1}{2} \sum_{i=0}^{w-1} \binom{w}{i} \sum_{j=0}^{i-1} \binom{n}{j} \right) \right)$.

Now we study the time and space complexity for steps 4 and 5 in both probabilistic and deterministic version. To represent each equation for the system of equation one needs $\sum_{w=0}^d \binom{n}{w}$ memory bits.

First we consider the probabilistic one. For probabilistic case one has to choose approximately $\frac{1}{2} \sum_{w=0}^d \binom{n}{w}$ many support input vectors of weight $> d$. Hence each linear equation obtained from these vectors has atleast $\sum_{i=0}^d \binom{d+1}{i}$ many coefficients of g and half of them can be eliminated using the equations obtained in previous steps. So, to get each equation one needs atleast $\sum_{w=0}^d \binom{n}{w} + \frac{1}{2} \sum_{i=0}^d ((\binom{d+1}{i} \sum_{j=0}^{i-1} \binom{n}{j}))$ computations. Hence the space complexity during 4th step is $SP2 \geq \frac{1}{2} (\sum_{w=0}^d \binom{n}{w})^2$ and time complexity is $TP2 \geq \frac{1}{2} \sum_{w=0}^d \binom{n}{w} (\sum_{w=0}^d \binom{n}{w} + \frac{1}{2} \sum_{i=0}^d ((\binom{d+1}{i} \sum_{j=0}^{i-1} \binom{n}{j})))$. Finally, to generate system of homogenous linear equations one requires

$SP = S1 + SP2 \geq \frac{1}{2} \sum_{w=0}^d ((\binom{n}{w} \sum_{i=0}^w \binom{n}{i})) + \frac{1}{2} (\sum_{w=0}^d \binom{n}{w})^2$ memory bits and
 $TP = T1 + TP2 \geq \frac{1}{2} \sum_{w=0}^d ((\binom{n}{w} (\sum_{i=0}^w \binom{n}{i} + \frac{1}{2} \sum_{i=0}^{w-1} \binom{w}{i} \sum_{j=0}^{i-1} \binom{n}{j})))$
 $+ \frac{1}{2} \sum_{w=0}^d \binom{n}{w} (\sum_{w=0}^d \binom{n}{w} + \frac{1}{2} \sum_{i=0}^d ((\binom{d+1}{i} \sum_{j=0}^{i-1} \binom{n}{j})))$ computations. Then in step 5, we have to solve $\frac{1}{2} \sum_{w=0}^d \binom{n}{w}$ many linear equations with same number of variables. To solve this system one needs $TP3 = (\frac{1}{2} \sum_{w=0}^d \binom{n}{w})^3$ computations using the Gaussian elimination technique.

Now we study space and time complexity for deterministic one. Since f is balanced, there are approximately $2^{n-1} - \frac{1}{2} \sum_{w=0}^d \binom{n}{w} = \frac{1}{2} \sum_{w=d+1}^n \binom{n}{w}$ many support vectors having weight $> d$ and these many are considered to find out equations. Hence each linear equation obtained from these vectors of weight $w > d$ contains $\sum_{i=0}^d \binom{w}{i}$ many coefficients of g and half of them can be eliminated using the equations obtained in steps 1, 2 and 3. To get this equation one needs $\sum_{i=0}^d \binom{n}{i} + \frac{1}{4} \sum_{i=0}^d ((\binom{w}{i} \sum_{j=0}^{i-1} \binom{n}{j}))$ computations. Hence the total space complexity during 4th step is $SD2 = \frac{1}{4} \sum_{w=d+1}^n \binom{n}{w} \sum_{w=0}^d \binom{n}{d}$ and time complexity is $TD2 = \frac{1}{2} \sum_{w=d+1}^n \binom{n}{w} (\sum_{i=0}^d \binom{n}{i} + \frac{1}{4} \sum_{i=0}^d ((\binom{w}{i} \sum_{j=0}^{i-1} \binom{n}{j})))$. Finally, to generate homogenous linear equations one needs

$SD = S1 + SD2 = \frac{1}{2} \sum_{w=0}^d ((\binom{n}{w} \sum_{i=0}^w \binom{n}{i})) + \frac{1}{4} \sum_{w=d+1}^n \binom{n}{w} \sum_{w=0}^d \binom{n}{d}$ memory bits and
 $TD = T1 + TD2 = \frac{1}{2} \sum_{w=0}^d ((\binom{n}{w} (\sum_{i=0}^w \binom{n}{i} + \frac{1}{2} \sum_{i=0}^{w-1} \binom{w}{i} \sum_{j=0}^{i-1} \binom{n}{j})))$
 $+ \frac{1}{2} \sum_{w=d+1}^n \binom{n}{w} (\sum_{i=0}^d \binom{n}{i} + \frac{1}{4} \sum_{i=0}^d ((\binom{w}{i} \sum_{j=0}^{i-1} \binom{n}{j})))$ computations. Further, in step 5, we have to solve $\frac{1}{2} \sum_{w=d+1}^n \binom{n}{w}$ many linear equations with $\frac{1}{2} \sum_{w=0}^d \binom{n}{w}$ number of variables. To solve this system one needs $TD3 = (\frac{1}{2} \sum_{w=d+1}^n \binom{n}{w})^3$ computations.

The system of equations generated by our strategy as well as Meier et al [23] algorithms are same. So, it takes same complexities to solve them. Only difference is during generation of the system of equations. In the following table we show the complexities for both algorithms for generating the system of equations.

4 Further reduction in matrix size applying linear transformation over the input variables of the function

To check for the annihilators, we need to compute the rank of the matrix UA^r . Following Theorem 3, it is clear that the size of the matrix UA^r will decrease

	Space	Time
Meier's algorithm	$\frac{1}{2} \sum_{w=0}^d \binom{n}{w} \sum_{i=0}^w \binom{n}{i}$ $+ \frac{1}{2} (\sum_{w=0}^d \binom{n}{w})^2$	$\frac{1}{2} \sum_{w=0}^d \binom{n}{w} (\sum_{i=0}^w \binom{n}{i}) + \frac{1}{2} \sum_{i=0}^{w-1} \binom{w}{i} \sum_{j=0}^{i-1} \binom{n}{j}$ $+ \frac{1}{2} \sum_{w=0}^d \binom{n}{w} (\sum_{i=0}^d \binom{n}{i}) + \frac{1}{2} \sum_{i=0}^d \binom{d+1}{i} \sum_{j=0}^{i-1} \binom{n}{j}$
Our algorithm	$\frac{1}{4} (\sum_{w=0}^d \binom{n}{w})^2$	$\frac{1}{4} (\sum_{w=0}^d \binom{n}{w})^2$

Table 1. Time and Space complexity comparison of Probabilistic algorithms to generate equations.

	Space	Time
Meier's algorithm	$\frac{1}{2} \sum_{w=0}^d \binom{n}{w} \sum_{i=0}^w \binom{n}{i}$ $+ \frac{1}{4} \sum_{w=d+1}^n \binom{n}{w} \sum_{i=0}^d \binom{n}{i}$	$\frac{1}{2} \sum_{w=0}^d \binom{n}{w} (\sum_{i=0}^w \binom{n}{i}) + \frac{1}{2} \sum_{i=0}^{w-1} \binom{w}{i} \sum_{j=0}^{i-1} \binom{n}{j}$ $+ \frac{1}{2} \sum_{w=d+1}^n \binom{n}{w} (\sum_{i=0}^d \binom{n}{i}) + \frac{1}{4} \sum_{i=0}^d \binom{w}{i} \sum_{j=0}^{i-1} \binom{n}{j}$
Our algorithm	$\frac{1}{4} \sum_{w=d+1}^n \binom{n}{w} \sum_{i=0}^d \binom{n}{i}$	$\frac{1}{4} \sum_{w=d+1}^n \binom{n}{w} \sum_{i=0}^d \binom{n}{i}$

Table 2. Time and Space complexity comparison of Deterministic algorithms to generate equations.

if μ_f increases and ν_f decreases. Let B be an $n \times n$ nonsingular binary matrix and b be an n -bit vector. The function $f(x)$ has an annihilator at degree d iff $f(Bx + b)$ has an annihilator at degree d . Thus one will try to get the affine transformation on the input variables of $f(x)$ to get $h(x) = f(Bx + b)$ such that $|\{x | h(x) = 1, wt(x) \leq d\}|$ is maximized. This is because in this case μ_h will be maximized and ν_h will be minimized and hence the dimension of the matrix UA^r , i.e., $\nu_f \times (\sum_{i=0}^d \binom{n}{i} - \mu_f)$ will be minimized. This will indeed decrease the complexity at the construction step (discussed in the previous section). More importantly, it will decrease the complexity to solve the system of homogeneous linear equations.

See the following example that explains the efficiency for a 5-variable function.

Example 2. We present an example for this purpose. Consider the 5-variable Boolean function f constructed using the method presented in [18] such that neither f nor $1 + f$ has an annihilator at a degree < 3 . The standard truth table representation of the function is 01010110010101100101001101001, i.e., the outputs are corresponding to the inputs which are of increasing value. One can check that $|\{x \in \{0, 1\}^5 \mid f(x) = 1 \text{ \& } wt(x) < 3\}| = 6$. Now if we consider the

function $h(x) = f(Bx + b)$ such that $B = \begin{bmatrix} 1 & 1 & 1 & 0 & 1 \\ 1 & 1 & 1 & 1 & 0 \\ 1 & 0 & 1 & 0 & 0 \\ 1 & 1 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 \end{bmatrix}$, and $b = \{1, 1, 0, 0, 1\}$, then

$|\{x \in \{0, 1\}^5 \mid h(x) = 1 \text{ \& } wt(x) < 3\}| = 16$ and one can immediately conclude (from the results in [19]) that neither h nor $1 + h$ has an annihilator of degree < 3 . This is an example where after finding the affine transformation there is even no need for the solution step at all. For the function f , here $h(x) = f(Bx + b)$ such that $|\{x | h(x) = 1, wt(x) \leq d\}|$ is maximized.

We also present an example for a sub optimal case. In this case we consider $B = \begin{bmatrix} 1 & 0 & 1 & 0 & 0 \\ 1 & 1 & 0 & 0 & 0 \\ 1 & 1 & 1 & 0 & 1 \\ 0 & 0 & 0 & 1 & 1 \\ 0 & 1 & 1 & 1 & 0 \end{bmatrix}$, and b an all zero vector, then $|\{x \in \{0,1\}^5 \mid h(x) = 1 \ \& \ wt(x) < 3\}| = 14$. Thus the dimension of the matrix UA^r becomes 2×2 as $\nu_f = 2$ and $\sum_{i=0}^d \binom{n}{i} - \mu_f = 2$. Thus one needs to check the rank of a 2×2 matrix only.

Now the question is how to find such an affine transformation (for the optimal or even for sub optimal cases) efficiently.

For exhaustive search to get the optimal affine transform one needs to check $f(Bx + b)$ for all $n \times n$ nonsingular binary matrices B and n bit vectors b . Since there are $\prod_{i=0}^{n-1} (2^n - 2^i)$ many nonsingular binary matrices and 2^n many n bit vectors, one needs to check $2^n \prod_{i=0}^{n-1} (2^n - 2^i)$ many cases for an exhaustive search. As weight of the input vectors are invariant under permutation of the arguments, checking for only one nonsingular matrix from the set of all nonsingular matrices whose rows are equivalent under certain permutation will suffice. Hence the exact number of search options is $\frac{1}{n!} 2^n \prod_{i=0}^{n-1} (2^n - 2^i)$. One can check for $n \times n$ nonsingular binary matrices B where $row_i < row_j$ for $i < j$ (row_i is the decimal value of binary pattern of i th row). It is clear that the search is infeasible for $n \geq 8$.

Now we present a heuristic towards this. Our aim is to find out an affine transformation $h(x)$ of $f(x)$, i.e., $h(x) = f(Bx + b)$, which maximizes the value of μ_h . This means the weight of the most of the input vectors having weight $\leq d$ should be in $supp(h)$. So we attempt to get an affine transformation for a Boolean function f such that the transformation increases the probability that an input vector, having output 1, will be translated to a low weight input vector.

Consider $h(Vx + v) = f(x)$, where V is an $n \times n$ binary matrix and $v = (v_1, v_2, \dots, v_n) \in \{0,1\}^n$. Suppose $r_1, r_2, \dots, r_n \in \{0,1\}^n$ are the row vectors of the transformation V . By $Vx + v = y$ we mean $Vx^{tr} + v = y^{tr}$, where $x = (x_1, x_2, \dots, x_n), y = (y_1, y_2, \dots, y_n) \in \{0,1\}^n$. Given an x , we find a y by this transformation and then $h(y)$ is assigned to the value of $f(x)$. If $f(x) = 1$, we like that the corresponding $y = Vx + v$ should be of low weight. The chance of (y_1, y_2, \dots, y_n) getting low weight increases if the probability of $y_i = 0, 1 \leq i \leq n$ is increased. That means the probability of $r_i \cdot (x_1, x_2, \dots, x_n) + v_i = 0$ for $1 \leq i \leq n$ needs to be increased. Hence we will like to choose a linearly independent set $r_i \in \{0,1\}^n, 1 \leq i \leq n$ and $v \in \{0,1\}^n$ such that the probability $r_i \cdot (x_1, x_2, \dots, x_n) + b_i = 0, 1 \leq i \leq n$ is high when $(x_1, x_2, \dots, x_n) \in supp(f)$. Since we use the relations $h(Vx + v) = f(x)$, and $h(x) = f(Bx + b)$, that means $B = V^{-1}$ and $b = V^{-1}v$.

The heuristic is presented below. By $bin[i]$ we denote the n -bit binary representation of the integer i .

Heuristic 1

1. $loop = 0; max = |\{x | f(x) = 1, wt(x) \leq d\}|;$
2. For $(i = 1; i < 2^n; i++)$ {
 - (a) $t = |\{x = (x_1, x_2, \dots, x_n) \in supp(f) | bin[i] \cdot x = 0\}|$
 - (b) if $t \geq \frac{wt(f)}{2}$, $val[i] = t$ and $a_i = 0$ else $val[i] = wt(f) - t$ and $a_i = 1$.
3. Arrange the triplets $(bin[i], a_i, val[i])$ in descending order of $val[i]$.
4. Choose suitable n many triplets (r_j, v_j, k_j) for $1 \leq j \leq n$ such that r_j 's are linearly independent and k_j 's are high.
5. Construct the nonsingular matrix V taking r_j , $1 \leq j \leq n$ as j -th row and $v = (v_1, v_2, \dots, v_n)$.
6. Increment $loop$ by 1; while $(loop < maxval)$
 - (a) $B = V^{-1}$, $b = V^{-1}v$.
 - (b) if $max < |\{x | f(Bx + b) = 1, wt(x) \leq d\}|$ replace $f(x)$ by $f(Bx + b)$ and update max by $|\{x | f(Bx + b) = 1, wt(x) \leq d\}|$.
 - (c) Go to step 2.

The time complexity of this heuristic is $(maxval \times n2^{2n})$. See the following example, where we trace Heuristic 1 for the 5-variable function f given in Example 2.

Example 3. We have $f = 01010110010101100101011001101001$ and check that $|\{x \in \{0,1\}^5 | f(x) = 1 \ \& \ wt(x) \leq 2\}| = 6$. In step 2, we get $(val[i], a_i)$ for $1 \leq i \leq 31$ as 1 : (11, 1), 2 : (8, 1), 3 : (11, 1), 4 : (8, 1), 5 : (11, 1), 6 : (8, 1), 7 : (9, 0), 8 : (8, 1), 9 : (9, 1), 10 : (8, 1), 11 : (9, 1), 12 : (8, 1), 13 : (9, 1), 14 : (8, 1), 15 : (11, 0), 16 : (8, 1), 17 : (9, 1), 18 : (8, 1), 19 : (9, 1), 20 : (8, 1), 21 : (9, 1), 22 : (8, 1), 23 : (11, 0), 24 : (8, 1), 25 : (9, 0), 26 : (8, 1), 27 : (9, 0), 28 : (8, 1), 29 : (9, 0), 30 : (8, 1), 31 : (11, 1). Then after ordering according the value of $val[i]$, we choose the row of matrix V as the 5-bit binary expansion of 1, 3, 5, 15 and 7 with frequency values of 0's as 11, 11, 11, 11, 9 respectively and $v = (a_1, a_3, a_5, a_{15}, a_7) = (1, 1, 1, 1, 0)$. Here the matrix V is a nonsingular matrix. The new function is $g = f(Bx + b)$, where $B = V^{-1}$, $b = V^{-1}v$ and one can check that $|\{x \in \{0,1\}^5 | g(x) = 1 \ \& \ wt(x) \leq 2\}| = 16$.

Experiments with this heuristic on different Boolean functions provide very positive results. First of all we have considered the functions which are random affine transformations $g(x)$ of the function [19], $f_s(x) = 1$ for $wt(x) \leq \lfloor \frac{n-1}{2} \rfloor$ and $f_s(x) = 0$ for $wt(x) \geq \lfloor \frac{n+1}{2} \rfloor$, which has no annihilator having degree $\leq \lfloor \frac{n-1}{2} \rfloor$. This experimentation has been done for $n = 5$ to 16. For all the cases running Heuristic 1 on $g(x)$ we could go back to $f_s(x)$. Then we have randomly changed $2^{\zeta n}$ bits on the upper half of $f_s(x)$ ($0.5 \leq \zeta \leq 0.8$ at steps of 0.1) to get $f'_s(x)$ and then put random transformations on $f'_s(x)$ to get $g(x)$. Running Heuristic 1, we could also go back to $f'_s(x)$ easily. For experiments we have taken $maxval = 20$.

The important issue is exactly when this matrix size is asymptotically reduced than the trivial matrix size $wt(f) \times \sum_{i=0}^d \binom{n}{i}$ if one writes down the equations by looking at the truth table of the function only. This happens only when μ_f is very close to $\sum_{i=0}^d \binom{n}{i}$. Let $\sum_{i=0}^d \binom{n}{i} - \mu_f \leq 2^{\zeta n}$, where ζ is a constant such that $0 < \zeta < 1$. In that case the matrix size will be less than or equal

to $(wt(f) + 2^{\zeta n} - \sum_{i=0}^d \binom{n}{i}) \times 2^{\zeta n}$. When $d = \lfloor \frac{n}{2} \rfloor$ and n odd, $\sum_{i=0}^d \binom{n}{i} = 2^{n-1}$. Thus for a balanced function, the size of the matrix becomes as low as $2^{\zeta n} \times 2^{\zeta n}$. We summarize the result as follows.

Theorem 4. *Predetermine a constant ζ , such that $0 < \zeta < 1$. Consider any Boolean function $f(x) \in B_n$ for which there exist a nonsingular binary matrix B and an n -bit vector b such that $\sum_{i=0}^d \binom{n}{i} - |\{x | f(Bx + b) = 1, wt(x) \leq d\}| \leq 2^{\zeta n}$. If B and b are known, then the size of the matrix UA^r will be less than or equal to $(wt(f) + 2^{\zeta n} - \sum_{i=0}^d \binom{n}{i}) \times 2^{\zeta n}$ which is asymptotically reduced in size than $wt(f) \times \sum_{i=0}^d \binom{n}{i}$.*

That B, b can be known is quite likely from the experimental results available running Heuristic 1.

Next we have run our heuristics on randomly chosen balanced functions. The number of inputs up to weight d for a Boolean function is $\sum_{i=0}^d \binom{n}{i}$. Thus for a randomly chosen balanced function, it is expected that there will be $\frac{1}{2} \sum_{i=0}^d \binom{n}{i}$ many inputs up to weight d for which the outputs are 1. Below we present the improvement (on an average of 100 experiments in each case) we got after running Heuristic 1 with $maxval = 20$ for $n = 12$ to 16.

n	12			13			14			15			16		
d	3	4	5	4	5	6	4	5	6	5	6	7	5	6	7
$\sum_{i=0}^d \binom{n}{i}$	299	794	1586	1093	2380	4096	1471	3473	6476	4944	9949	16384	6885	14893	26333
$\lceil \frac{1}{2} \sum_{i=0}^d \binom{n}{i} \rceil$	149	397	793	541	1190	2048	735	1736	3238	2472	4974	8192	3442	7446	13166
Heuristic Value	228	535	964	717	1438	2322	957	2051	3648	2917	5525	8811	3995	8194	14114

Table 3. Efficiency of Heuristic 1 on random balanced functions.

It should be noted that after running our heuristic on random balanced functions, the improvement is not extremely significant. There are improvements as we find that the values are significantly more than $\frac{1}{2} \sum_{i=0}^d \binom{n}{i}$ (making our algorithm efficient), but the value is not very close to $\sum_{i=0}^d \binom{n}{i}$. This is not a problem with the efficiency of the heuristic, but with the inherent property of a random Boolean function that there may not be an affine transformation at all on $f(x)$ such that $|\{x | f(Bx + b) = 1, wt(x) \leq d\}|$ is very high. In fact we can show that for highly nonlinear functions $f(x)$, the increment from $|\{x | f(x) = 1, wt(x) \leq d\}|$ to $|\{x | f(Bx + b) = 1, wt(x) \leq d\}|$ may not be significant for any B, b . The reason for this is as follows.

Proposition 2. *Let $f \in B_n$ be a balanced function (n odd) having nonlinearity $nl(f) = 2^{n-1} - 2^{\frac{n-1}{2}}$. Then for any nonsingular $n \times n$ matrix B and any n -bit vector b , $2^{n-1} - |\{x | f(Bx + b) = 1, wt(x) \leq \frac{n-1}{2}\}| \geq \frac{1}{2} \binom{n-1}{\frac{n-1}{2}} - 2^{\frac{n-1}{2}-1}$.*

Proof. Let $f \in B_n$ be a balanced function (n odd) having nonlinearity $nl(f) = 2^{n-1} - 2^{\frac{n-1}{2}}$. Let $g \in B_n$ be the function such that $g(x) = 1$ for $wt(x) \leq \frac{n-1}{2}$. By [19, Theorem 3], $nl(g) = 2^{n-1} - \binom{n-1}{\frac{n-1}{2}}$. Now we like to find out a function

$h(x) = f(Bx + b)$ such that $|\{x|h(x) = 1, wt(x) \leq \frac{n-1}{2}\}|$ is high. Consider the value $T = |supp(g) \cap supp(h)|$, i.e., $T = |\{x : h(x) = 1 \ \& \ wt(x) \leq \frac{n-1}{2}\}|$. Without loss of generality consider $T \geq 2^{n-2}$. Hence, $d(h, g) = 2(2^{n-1} - T) = 2^n - 2T$. Now, $nl(f) = nl(h) \leq nl(g) + d(h, g) = (2^{n-1} - \binom{n-1}{\frac{n-1}{2}}) + 2^n - 2T$. Thus, $2^{n-1} - 2^{\frac{n-1}{2}} \leq (2^{n-1} - \binom{n-1}{\frac{n-1}{2}}) + 2^n - 2T$, i.e., $2^{n-1} - T \geq \frac{1}{2}(\binom{n-1}{\frac{n-1}{2}}) - 2^{\frac{n-1}{2}-1}$. \square

Thus if one predetermines a ζ , then for a large n we may not satisfy the condition that $\sum_{i=0}^{\frac{n-1}{2}} \binom{n}{i} - |\{x|f(Bx + b) = 1, wt(x) \leq d\}| \leq 2^{\zeta n}$. In this direction we present the following general result where the constraint of nonlinearity is removed.

Theorem 5. Suppose $f \in B_n$ be a randomly chosen balanced function. Then the probability to get an affine transformation such that

$$|\{x|f(Bx + b) = 1, wt(x) \leq \lfloor \frac{n-1}{2} \rfloor\}| > \sum_{i=0}^{\lfloor \frac{n-1}{2} \rfloor} \binom{n}{i} - k \text{ is}$$

1. less than $\frac{(n+1)2^n \sum_{i=0}^{k-1} \binom{2^{n-1}}{i}^2}{\binom{2^n}{2^{n-1}-1}}$ for n odd.
2. less than $\frac{(n+1)2^n \sum_{i=0}^{k-1} \left(\sum_{j=0}^{\frac{n}{2}-1} \binom{n}{j} \right) \binom{2^n - \sum_{j=0}^{\frac{n}{2}-1} \binom{n}{j}}{i + \frac{1}{2} \binom{n}{\frac{n}{2}}} }{\binom{2^n}{2^{n-1}-1}}$ for n even.

Proof. First we prove it for n odd. The number of balanced functions $h \in B_n$ such that $|\{x|h(x) = 1, wt(x) \leq \frac{n-1}{2}\}| > 2^{n-1} - k$ is $\sum_{i=0}^{k-1} \binom{2^{n-1}}{i}^2$ (consider the upper and lower half in the truth table of the function). So, there will be at most $\sum_{i=0}^{k-1} \binom{2^{n-1}}{i}^2$ many affinely invariant classes of such functions. Further the total number of balanced function is $\binom{2^n}{2^{n-1}}$. Hence the total number of affinely invariant classes of balanced function is $\geq \frac{\binom{2^n}{2^{n-1}-1}}{2^n(2^n-1)(2^n-2^1)\dots(2^n-2^{n-1})} > \frac{\binom{2^n}{2^{n-1}-1}}{(n+1)2^n}$. Hence the probability of a randomly chosen balanced function will be function type h is bounded by $\frac{(n+1)2^n \sum_{i=0}^{k-1} \binom{2^{n-1}}{i}^2}{\binom{2^n}{2^{n-1}-1}}$. Similarly, the case for n even can be proved. \square

If one takes $k \leq 2^{\frac{3}{4}n}$, then it can be checked easily that the probability decreases fast towards zero as n increases. Thus for a random balanced function f , the probability of getting an affine transformation (which generates the function h from f) such that $|\{x|f(Bx + b) = 1, wt(x) \leq \lfloor \frac{n-1}{2} \rfloor\}| > \sum_{i=0}^{\lfloor \frac{n-1}{2} \rfloor} \binom{n}{i} - 2^{\frac{3}{4}n}$ is almost improbable.

Thus when one randomly chosen balanced function is considered, using the strategy of considering the function after affine transformation, one can indeed reduce the matrix size by constant factor, but the reduction may not be significant in asymptotic terms when the annihilators at the degree of $\lfloor \frac{n-1}{2} \rfloor$ are considered for large n .

5 Conclusion

In this paper we study how to reduce the matrix size which is involved in finding the annihilators of a Boolean function. Our results show that considerable reduction in the size of the matrix is achievable. We identify the classes where it provides asymptotic improvement. We also note that for randomly chosen balanced functions, the improvement is rather constant than asymptotic. The reduction in matrix size helps in running the actual annihilator finding steps by Gaussian elimination method. Though our method is less efficient in general than the recently known efficient algorithms [1, 21] to find the annihilators, this work helps in theoretically understanding the structure of the matrix involved.

Acknowledgment: We like to acknowledge one of the anonymous reviewers for pointing out some problems in Heuristic 1 in the submitted version of this paper; this is corrected in this final version.

References

1. F. Armknecht, C. Carlet, P. Gaborit, S. Kuenzli, W. Meier and O. Ruatta. Efficient computation of algebraic immunity for algebraic and fast algebraic attacks. In *Eurocrypt 2006*.
2. F. Armknecht. Improving Fast Algebraic Attacks. In *FSE 2004*, number 3017 in Lecture Notes in Computer Science, pages 65–82. Springer Verlag, 2004.
3. G. Ars and J. Faugère. Algebraic Immunities of functions over finite fields. INRIA report, 2005.
4. L. M. Batten. Algebraic Attacks over $\text{GF}(q)$. In *Progress in Cryptology - INDOCRYPT 2004*, pages 84–91, number 3348, Lecture Notes in Computer Science, Springer-Verlag.
5. A. Botev and Y. Tarannikov. Lower bounds on algebraic immunity for recursive constructions of nonlinear filters. Preprint 2004.
6. A. Braeken and B. Praneel. Probabilistic algebraic attacks. In *10th IMA international conference on cryptography and coding*, 2005.
7. A. Braeken and B. Praneel. On the Algebraic Immunity of Symmetric Boolean Functions. In *Indocrypt 2005*, number 3797 in LNCS, pages 35–48. Springer Verlag, 2005. Also available at Cryptology ePrint Archive, <http://eprint.iacr.org/>, No. 2005/245, 26 July, 2005.
8. A. Braeken, J. Lano and B. Praneel. Evaluating the Resistance of Stream Ciphers with Linear Feedback Against Fast Algebraic Attacks. Accepted in *ACISP 2006*.
9. A. Canteaut. Open problems related to algebraic attacks on stream ciphers. In *WCC 2005*, pages 1–10, invited talk.
10. C. Carlet. Improving the algebraic immunity of resilient and nonlinear functions and constructing bent functions. IACR ePrint server, <http://eprint.iacr.org>, 2004/276.
11. J. H. Cheon and D. H. Lee. Resistance of S-boxes against Algebraic Attacks. In *FSE 2004*, number 3017 in Lecture Notes in Computer Science, pages 83–94. Springer Verlag, 2004.
12. J. Y. Cho and J. Pieprzyk. Algebraic Attacks on SOBER-t32 and SOBER-128. In *FSE 2004*, number 3017 in Lecture Notes in Computer Science, pages 49–64. Springer Verlag, 2004.

13. N. Courtois and J. Pieprzyk. Cryptanalysis of block ciphers with overdefined systems of equations. In *Advances in Cryptology - ASIACRYPT 2002*, number 2501 in Lecture Notes in Computer Science, pages 267–287. Springer Verlag, 2002.
14. N. Courtois and W. Meier. Algebraic attacks on stream ciphers with linear feedback. In *Advances in Cryptology - EUROCRYPT 2003*, number 2656 in Lecture Notes in Computer Science, pages 345–359. Springer Verlag, 2003.
15. N. Courtois. Fast algebraic attacks on stream ciphers with linear feedback. In *Advances in Cryptology - CRYPTO 2003*, number 2729 in Lecture Notes in Computer Science, pages 176–194. Springer Verlag, 2003.
16. N. Courtois, B. Debraize and E. Garrido. On Exact Algebraic [Non-]Immunity of S-boxes Based on Power Functions. Accepted in *ACISP 2006*.
17. D. K. Dalai, K. C. Gupta and S. Maitra. Results on Algebraic Immunity for Cryptographically Significant Boolean Functions. In *INDOCRYPT 2004*, pages 92–106, number 3348, Lecture Notes in Computer Science, Springer-Verlag.
18. D. K. Dalai, K. C. Gupta and S. Maitra. Cryptographically Significant Boolean functions: Construction and Analysis in terms of Algebraic Immunity. In *FSE 2005*, pages 98–111, number 3557, Lecture Notes in Computer Science, Springer-Verlag.
19. D. K. Dalai, S. Maitra and S. Sarkar. Basic Theory in Construction of Boolean Functions with Maximum Possible Annihilator Immunity. Cryptology ePrint Archive, <http://eprint.iacr.org/>, No. 2005/229, 15 July, 2005.
20. D. K. Dalai and S. Maitra. Towards an Efficient Algorithm to find Annihilators by Solving a Set of Homogeneous Linear Equations. Cryptology ePrint Archive, <http://eprint.iacr.org/>, No. 2006/032, January 2006.
21. F. Didier and J. Tillich. Computing the Algebraic Immunity Efficiently. In *FSE 2006*.
22. D. H. Lee, J. Kim, J. Hong, J. W. Han and D. Moon. Algebraic Attacks on Summation Generators. In *FSE 2004*, Lecture Notes in Computer Science, pages 34–48. Springer Verlag, 2004.
23. W. Meier, E. Pasalic and C. Carlet. Algebraic attacks and decomposition of Boolean functions. In *Advances in Cryptology - EUROCRYPT 2004*, number 3027 in Lecture Notes in Computer Science, pages 474–491. Springer Verlag, 2004.
24. Y. Nawaz, G. Gong and K. C. Gupta. Upper Bounds on Algebraic Immunity of Power Functions. In *FSE 2006*.