

Some remarks on how to hash faster onto elliptic curves

Dmitrii Koshelev¹

Computer sciences and networks department, Télécom Paris

Abstract. In this article we propose three optimizations of indifferentiable hashing onto (prime order subgroups of) ordinary elliptic curves over finite fields \mathbb{F}_q . One of them is dedicated to elliptic curves E provided that $q \equiv 11 \pmod{12}$. The other two optimizations take place respectively for the subgroups $\mathbb{G}_1, \mathbb{G}_2$ of some pairing-friendly curves. The performance gain comes from the smaller number of required exponentiations in \mathbb{F}_q for hashing to $E(\mathbb{F}_q), \mathbb{G}_2$ (resp. from the absence of necessity to hash directly onto \mathbb{G}_1). In particular, our results affect the pairing-friendly curve BLS12-381 (the most popular in practice at the moment) and the (unique) French curve FRP256v1 as well as almost all Russian standardized curves and a few ones from the draft NIST SP 800-186.

Key words: BLS12 family of pairing-friendly curves, clearing cofactor, indifferentiable hashing to elliptic curves, optimal ate pairings.

1 How to hash onto pairing-friendly curves

This is an addendum to our recent articles [1], [2], [3]. So, with your permission, we do not provide a detailed introduction in order to avoid repetition. Good surveys on how to hash into (or onto) elliptic curves over finite fields are also represented in [4, §8], [5]. It is worth emphasizing that throughout this text we mean hashing indifferentiable from a random oracle (in the sense of [12, §2.2]).

Let E_1 be an ordinary pairing-friendly elliptic curve of embedding degree $k > 1$ over a finite field \mathbb{F}_q . Besides, let E_2 be a twist of E_1 of degree $d := \#\text{Aut}(E_1)$ over the field \mathbb{F}_{q^e} , where $e := k/d \in \mathbb{N}$. As is customary, for a common prime divisor r of the orders $N_1 := \#E_1(\mathbb{F}_q)$ and $N_2 := \#E_2(\mathbb{F}_{q^e})$ denote by $\mathbb{G}_1 \subset E_1(\mathbb{F}_q)$ and $\mathbb{G}_2 \hookrightarrow E_2(\mathbb{F}_{q^e})$ the eigenspaces of the Frobenius endomorphism on $E_1[r] \subset E_1(\mathbb{F}_{q^k})$, associated with the eigenvalues 1, q respectively. Note that the condition $e \in \mathbb{N}$ is not automatically met, i.e., this is our assumption. It is claimed (e.g., in [4, Theorem 3.3.5]) that for any prime divisor $r \mid N_1$ there is always a unique non-trivial \mathbb{F}_{q^e} -twist E_2 (of degree d) such that $r \mid N_2$. By abuse of notation, we identify the order r subgroup $\mathbb{G}_2 \subset E_1(\mathbb{F}_{q^k})$ with its image under an \mathbb{F}_{q^e} -isomorphism $E_1 \simeq E_2$. Thus $\mathbb{G}_1 = E_1(\mathbb{F}_q)[r]$ and $\mathbb{G}_2 = E_2(\mathbb{F}_{q^e})[r]$. Besides, $d \in \{2, 4, 6\}$ and $d = 2$ if and only if $j(E_i) \neq 0, 1728$ (respectively, $d = 4$ iff $j(E_i) = 1728$ and $d = 6$ iff $j(E_i) = 0$).

This section explains how to hash onto \mathbb{G}_2 more efficiently and why we do not need to hash directly onto \mathbb{G}_1 . In the first case, we significantly exploit the presence of clearing the cofactor $c_2 := N_2/r$. In the second one, on the contrary, clearing the cofactor $c_1 := N_1/r$ can be fully avoided. The fact is that *optimal ate pairings* $a: \mathbb{G}_2 \times \mathbb{G}_1 \rightarrow \mu_r \subset \mathbb{F}_{q^k}^*$ [4, Theorem 3.3.4] can

¹web page: <https://www.researchgate.net/profile/Dimitri-Koshelev>
email: dimitri.koshelev@gmail.com

be painlessly (unlike $E_2(\mathbb{F}_{q^e}) \times \mathbb{G}_1$) extended to $\mathbb{G}_2 \times E_1(\mathbb{F}_q)$, at least in main pairing-based protocols.

At the moment, due to [6, Table 1] the curve BLS12-381 is a de facto standard in pairing-based cryptography. More generally, the *Barreto–Lynn–Scott family* with $k = 12$ and $d = 6$ (see, e.g., [7, §3.1]) possesses the parameters

$$r(z) = z^4 - z^2 + 1, \quad q(z) = (z - 1)^2 r(z) / 3 + z.$$

By definition, BLS12-381 is generated by $z := -0xd201000000010000$ and hence

$$\lceil \log_2(-z) \rceil = 64, \quad \lceil \log_2(r) \rceil = 255, \quad \lceil \log_2(q) \rceil = 381.$$

Notice that $r \ll q$ in contrast to the *Barreto–Naehrig family* [4, Example 4.2].

Earlier for the curve BLS12-381 (and a number of others) the author constructed in [1], [2] encodings $h_1: \mathbb{F}_q^2 \rightarrow E_1(\mathbb{F}_q)$ and $h_2: \mathbb{F}_q \rightarrow E_2(\mathbb{F}_{q^2})$ computable in constant time of one exponentiation in \mathbb{F}_q . Combining the new ideas with these encodings, we obtain hash functions $\{0, 1\}^* \rightarrow E_1(\mathbb{F}_q)$ and $\{0, 1\}^* \rightarrow \mathbb{G}_2$, which seem to be difficult to speed up even more. We will discuss (respectively in §1.1, §1.2) how much performance gain they provide over hash functions that are actively applied in the industry.

Recall that almost all known hash functions $\mathcal{H}_i: \{0, 1\}^* \rightarrow \mathbb{G}_i$ are the compositions $\mathcal{H}_i = [c'_i] \circ h_i \circ \eta_i$. Here $\eta_i: \{0, 1\}^* \rightarrow S_i$ are hash functions to some finite sets, $h_1: S_1 \rightarrow E_1(\mathbb{F}_q)$ and $h_2: S_2 \rightarrow E_2(\mathbb{F}_{q^e})$ are just maps traditionally called *encodings*, and finally $c'_i \in \mathbb{N}$ such that $c_i \mid c'_i$, $r \nmid c'_i$. The scalar multiplication $[c'_i]$ on the curve E_i is said to be *clearing cofactor*. Surprisingly, due to Fuentes-Castaneda et al. [8] it is more efficient to multiply points by scalars c'_i greater than c_i . The sets S_i are usually very simple, hence it is easy to combine η_i from existing hash functions $\{0, 1\}^* \rightarrow \{0, 1\}^\ell$ for $\ell \in \mathbb{N}$. The most complicated component of \mathcal{H}_i is no doubt h_i , because its essence is based on high-dimensional algebraic geometry.

The majority of pairing-based protocols requires a hash function to at most one group \mathbb{G}_1 or \mathbb{G}_2 . Of course, any such protocol can be equivalently implemented for hashing to the other group. Without using point compression-decompression methods, elements of \mathbb{G}_1 (resp. \mathbb{G}_2) are obviously represented by $2\lceil \log_2(q) \rceil$ (resp. $2e\lceil \log_2(q) \rceil$) bits. Therefore the choice often depends on whether a hash value should be more compact than the second pairing argument or vice versa. Besides, there are rarely used protocols, for example the Scott identity-based key agreement [9], where both hash functions \mathcal{H}_i are necessary. Thus the more cumbersome hashing to \mathbb{G}_2 can not be replaced by hashing to \mathbb{G}_1 in all situations.

1.1 How not to hash onto \mathbb{G}_1

As far as we know, (non-degenerate) optimal ate pairings $a: \mathbb{G}_2 \times \mathbb{G}_1 \rightarrow \mu_r \subset \mathbb{F}_{q^k}^*$ are only used in today's real-world cryptography. The fact is that the corresponding Miller loop has the hypothetically smallest length $\approx \log_2(r) / \varphi(k)$, where φ is Euler's totient function. However it is more practical to take the whole group $E_1(\mathbb{F}_q)$ instead of \mathbb{G}_1 . In this case, the pairing $a: \mathbb{G}_2 \times E_1(\mathbb{F}_q) \rightarrow \mu_r$ becomes degenerate, but this is not important. A similar trick is done in [10, §5] for the Tate pairing in the context of isogeny-based cryptography, where, on the contrary, \mathbb{G}_2 is replaced by $E_1(\mathbb{F}_{q^k})$ in our notation.

Indeed, first, the length of the Miller loop depends only on the order of \mathbb{G}_2 . Second, if for points $P \in E_1(\mathbb{F}_q)$ and $Q \in \mathbb{G}_2$ we have $a(Q, P) = 1$, then a fortiori $a(Q, c'_1 P) = a(Q, P)^{c'_1} =$

1. We stress that popular protocols (such as the Boneh–Franklin identity-based encryption [4, §1.6.4] or the aggregated BLS signature [11]) work correctly whether the order of P equals r or not. Nevertheless, it should be borne in mind that the *strong unforgeability property* (unlike the usual *existential* one) is not satisfied anymore as emphasized in [11, §5.2]. Finally, the complexity of computing $a(Q, P)$ remains the same as that of computing $a(Q, c'_1 P)$, because $P, c'_1 P$ are equally defined over \mathbb{F}_q .

In [1] we construct an encoding $h_1: \mathbb{F}_q^2 \rightarrow E_1(\mathbb{F}_q)$ for elliptic curves $E_1: y^2 = x^3 + b$ (of j -invariant 0) provided that $\sqrt{b} \in \mathbb{F}_q$. There we prove that h_1 is *admissible* in the sense of [12, Definition 4], which leads (in compliance with [12, Theorem 1]) to the indifferentiable hash function $h_1 \circ \eta_1$. Moreover, h_1 can be implemented in constant time of raising to some power $n_1 \in \mathbb{N}$ in the field \mathbb{F}_q (not counting a few additions and multiplications). In particular, our encoding is applicable to the curve BLS12-381 for which $b = 4$ and $n_1 = (q - 10)/27$.

Recall that the famous (*indirect*) *Wahby–Boneh encoding* h_{WB} [13, §4] (based on the *simplified SWU* one [12, §7]) is also valid for BLS12-381. It requires to extract one square root in \mathbb{F}_q , which for that curve is equivalent to raising in \mathbb{F}_q to the power $n_2 := (q - 3)/4 \in \mathbb{N}$. The hash function H_2 from [13, §5] twice applies h_{WB} in order to act as a random oracle. By the way, the other indifferentiable hash function H_3 is even slower than H_2 by virtue of [13, Figure 1].

To be exact, the Hamming weight $w(n_1) = 192$ and $w(n_2) = 228$. Denote by $\ell(n_i)$ the length of a shortest addition chain for n_i . In accordance with [14, §9.2.1] we obtain the inequalities

$$382 \leq \ell(n_1) \lesssim 419, \quad 385 \leq \ell(n_2) \lesssim 422.$$

We can not claim that these upper bounds are mathematically correct, because we omitted $o(1)$ in the original inequality. However, in any case, the sought bounds are very close (probably equal) to ours.

On the other hand, following the sliding window method [14, §9.1.3] (with $k = 5$), one can explicitly derive an addition chain for n_1 (resp. n_2) whose the length equals 449 (resp. 458). We invite the reader to independently check our conclusion, since the mentioned method is simple and has many public implementations. Curiously, a similar chain for n_2 of the same length 458, obtained by means of more advanced methods, appears in the optimized library [15]. Thus the encoding h_{WB} applied twice is much slower than ours h_1 applied once. Indeed, $2 \cdot 458 - 449 = 467$ is a significant amount of multiplications in \mathbb{F}_q that can be eliminated by giving priority to h_1 .

1.2 How to hash onto \mathbb{G}_2

To our knowledge, optimal ate pairings do not have a natural extension to $E_2(\mathbb{F}_{q^e}) \times \mathbb{G}_1$. Conversely, (non-degenerate) *twisted optimal ate pairings* [4, Theorem 3.3.8] of the form $\mathbb{G}_1 \times \mathbb{G}_2 \rightarrow \mu_r$ are readily extended to $\mathbb{G}_1 \times E_2(\mathbb{F}_{q^e})$. But, unfortunately, for them the Miller loop is of a larger length than for (usual) optimal ate pairings. It is generally recognized that a pairing is a more laborious operation than an elliptic curve scalar multiplication. Therefore reducing the Miller loop seems a better solution than avoiding the multiplication by c'_2 .

For the next theorem we need the notions of (*B*-) *well-distributed* [16, Definitions 5, 7] and (ϵ -) *regular map* [16, Definition 3] (with respect to the uniform distribution on its domain).

In the definition of an admissible map the regularity is essentially the only property that is difficult to satisfy. So it is necessary to focus on establishing it. For the sake of convenience, consider so-called *tensor multiplication* of any two maps $h: S \rightarrow G$, $g: T \rightarrow G$ from sets S , T to the same group $(G, +)$:

$$h \otimes g: S \times T \rightarrow G \quad (s, t) \mapsto h(s) + g(t).$$

Theorem 1. *Assume there is a B -well-distributed encoding $h_2: \mathbb{F}_q \rightarrow E_2(\mathbb{F}_{q^e})$ (for $B \in \mathbb{R}_{>0}$) and a point of $E_2(\mathbb{F}_{q^e})$ of order $m \mid c_2$ (or, equivalently, $m \mid c'_2$). Then the map $[c'_2] \circ h_2: \mathbb{F}_q \rightarrow \mathbb{G}_2$ is ϵ -regular, where $\epsilon := B\sqrt{N_2/(mq)}$.*

Proof. Pick any point $P \in E_2(\mathbb{F}_{q^e})$ of order m . According to [16, Corollary 1] the map $F := h_2 \otimes g$, where

$$g: [0, m) \rightarrow E_2(\mathbb{F}_{q^e}) \quad t \mapsto tP,$$

is ϵ -regular for ϵ as in the statement of the theorem. It is readily checked that the composition $[c'_2] \circ F$ is still ϵ -regular. Since $[c'_2] \circ h_2 = [c'_2] \circ F$, the theorem is proved. \square

For $e = 2$ an example of the desired encoding h_2 is given in [2] (modulo notation) as the composition $h_2 := \psi \circ \varphi \circ h$. Here $h: \mathbb{F}_q \rightarrow H(\mathbb{F}_q)$ is an encoding to some \mathbb{F}_q -curve H of geometric genus two, $\varphi: H \rightarrow E'$ is a (quadratic) \mathbb{F}_{q^2} -cover to an auxiliary elliptic \mathbb{F}_{q^2} -curve E' of j -invariant $\notin \mathbb{F}_q$, and finally $\psi: E' \rightarrow E_2$ is an \mathbb{F}_{q^2} -isogeny of small degree. By virtue of [3, Corollary 1], [2, Theorem 1] the encodings h and $\varphi \circ h$ are 2-well-distributed. The same is true for h_2 whenever $\psi: E'(\mathbb{F}_{q^2}) \xrightarrow{\sim} E_2(\mathbb{F}_{q^2})$, which follows if $(\deg(\psi), N_2) = 1$.

For the BLS12 family we have the parametrizations

$$c_2(z) = (z^8 - 4z^7 + 5z^6 - 4z^4 + 6z^3 - 4z^2 - 4z + 13)/9, \quad c'_2(z) = 3(z^2 - 1)c_2(z)$$

according to [7, §4.1]. Recall that BLS12-381 has the form $E_2: y^2 = x^3 + 4(1 + i)$ (where $i := \sqrt{-1} \notin \mathbb{F}_q$) and, as mentioned in [2, Introduction], there is the desired isogeny ψ of degree $7 \nmid N_2$. Besides, the group $E_2(\mathbb{F}_{q^2})$ possesses a point of order $m = c_2/(13 \cdot 23)$, because this number is square free. As a result,

$$\epsilon = 2\sqrt{13 \cdot 23r/q} \leq 2^{-115/2}$$

is a negligible value. Incidentally, this can not be said about BN curves, since for them $r/q = 1 + O(q^{-1/2})$.

It is worth noting that the encoding h can be computed in constant time of extracting one square root in \mathbb{F}_q . This is equally true for h_2 , since φ, ψ are algebraic maps of small degrees. Among other things, the denominators of their defining functions do not need to be inverted, because *Jacobian projective coordinates* (see, e.g., [13, §2]) are preferred for use in practice.

By analogy with Theorem 1, the map $[c'_2] \circ \text{Map}_2$ (for Map_2 from [13, §5]) also turns out to be regular, that is the hash function H_4 from there actually acts as a random oracle. However this circumstance was not noticed in that article. In comparison to the Wahby–Boneh encoding h_{WB} , ours h_2 nevertheless allows to avoid extracting one square root in \mathbb{F}_q . The fact is that a square root in \mathbb{F}_{q^2} (which appears in the simplified SWU map), as is well known, can be expressed via two square roots in \mathbb{F}_q . By the way, the other hash functions H_5, H_6 are even slower than H_4 by virtue of [13, Figure 1].

2 How to hash onto $E(\mathbb{F}_q)$ if $q \equiv 11 \pmod{12}$

Hash functions to classical (i.e., non-pairing-friendly) elliptic curves have become more and more in demand. Indeed, according to [17, Table I] they are actively used in many PAKE (Password Authenticated Key Exchange) protocols. Several years ago CFRG (Crypto Forum Research Group) conducted the PAKE selection process [18] in which the protocols CPace [19] and OPAQUE [20] won. Besides, hashing to elliptic curve is necessary for some blind signatures (such as in [21, §3.3]), which serve as a basis, e.g., for electronic voting schemes.

Let us freely utilize notions arisen in previous sections. Consider an elliptic curve $E: y^2 = x^3 + ax + b$ defined over a finite field \mathbb{F}_q . Under the condition $q \equiv 2 \pmod{3}$ (resp. $j(E) \neq 0, 1728$), *Icart's encoding* h_I [22] (resp. the simplified SWU one h_{sSWU}) is available. In accordance with [22, Lemma 4], [12, Lemma 6] for any $P \in E(\mathbb{F}_q)$ we have $\#h_I^{-1}(P) \leq 4$ and $\#h_{sSWU}^{-1}(P) \leq 8$. In fact, if an implementation of h_{sSWU} takes into account the sign of the y -coordinate, then $\#h_{sSWU}^{-1}(P) \leq 4$. At the same time, by virtue of [23, §5] the encoding h_I (resp. h_{sSWU}) is B -well-distributed with $B = 13$ (resp. $B = 53$) at least for q of a cryptographic size. Applying [16, Corollary 1], we thus get

Theorem 2. *Suppose that $q \equiv 2 \pmod{3}$ and $j(E) \neq 0, 1728$. Then the map $F := h_I \otimes h_{sSWU}: \mathbb{F}_q^2 \rightarrow E(\mathbb{F}_q)$ is ϵ -regular for the negligible value $\epsilon := 26\sqrt{N}/q$, where $N := \#E(\mathbb{F}_q)$.*

From now on we assume in addition that $q \equiv 3 \pmod{4}$. Obviously,

$$q \equiv 2 \pmod{3}, q \equiv 3 \pmod{4} \quad \Leftrightarrow \quad q \equiv 11 \pmod{12}.$$

For the sake of compactness, we put $e := (q+1)/4$ and $k := (q+1)/12$. Notice that for $Z = n/d$ such that $n, d \in \mathbb{F}_q^*$ we obtain

$$z := Z^k = n^k \cdot d^{q-1-k} = n^k \cdot d^{(11q-13)/12} = nd^9 \cdot (nd^{11})^{(q-11)/12}, \quad z^6 = Z^{(q+1)/2} = \left(\frac{Z}{q}\right)Z,$$

where $\left(\frac{Z}{q}\right)$ is the Legendre symbol. In particular, $z = \sqrt[6]{Z}$ whenever Z is a quadratic residue in \mathbb{F}_q .

Given $(t, s) \in \mathbb{F}_q^2$ we need to evaluate $h_I(t)$ and $h_{sSWU}(s)$. As is known, separately each of these points can be computed in constant time of one exponentiation in \mathbb{F}_q (the case of h_{sSWU} see in [13, §4.2]). Let's show that this is also possible simultaneously for the two points (and hence for $F(t, s)$). The only cumbersome part of h_I (resp. h_{sSWU}) consists in the exponentiation $\sqrt[3]{f} = f^{(2q-1)/3}$ (resp. $\pm g^e$ such that $(g^e)^2 = \left(\frac{g}{q}\right)g$), where

$$f := \left(\frac{3a - t^4}{6t}\right)^2 - b - \frac{t^6}{27}, \quad g := -\frac{b}{a} \left(1 + \frac{1}{s^4 - s^2}\right).$$

Evidently, $\sqrt[3]{f}$ is the unique cubic root of f in \mathbb{F}_q and for our purpose it is sufficient to find g^e up to a sign. For the sake of simplicity, let us exclude from consideration the zeros and poles of the functions f, g . As usual, they can be processed individually.

We suggest to act in a similar way as in [24, §3], that is for $Z := f^2 g^3$ to compute $z = Z^k$ (almost $\sqrt[6]{Z}$) instead of separate computing $\sqrt[3]{f}$ and $\pm g^e$ (almost \sqrt{g}). Note that

$$z = f^{(q+1)/6} \cdot g^e = \left(\frac{f}{q}\right) \sqrt[3]{f} \cdot g^e, \quad z^2 = \sqrt[3]{f^2} \cdot \left(\frac{g}{q}\right)g.$$

Introducing the auxiliary notation $\theta := fg/z^2$, we get the equalities

$$\sqrt[3]{f} = \frac{\binom{g}{q}fg}{z^2} = \binom{g}{q}\theta, \quad g^e = \frac{z}{\binom{f}{q}\sqrt[3]{f}} = \frac{z}{\binom{fg}{q}\theta}.$$

We see that $\theta^3 = \binom{g}{q}f$ and $z^6 = \binom{g}{q}Z$. Therefore the symbol $\binom{g}{q}$ can be determined for free. More formally,

$$(\sqrt[3]{f}, \pm g^e) = \begin{cases} (\theta, z/\theta) & \text{if } \theta^3 = f, \text{ i.e., } z^6 = Z, \\ (-\theta, z/\theta) & \text{otherwise.} \end{cases}$$

Bearing in mind the formula above for $(n/d)^k$ without the inversion operation, we emphasize again that

Remark 1. *The map F (in contrast to $h_I^{\otimes 2}$ and $h_{sSWU}^{\otimes 2}$) can be computed in constant time of one exponentiation in \mathbb{F}_q .*

Of course, by analogy with §1.1, given q it is not difficult to derive explicit short addition chains for raising to the power k .

Since [12, Algorithm 1] continues to be correct for two different encodings, we eventually establish

Corollary 1. *The map $F: \mathbb{F}_q^2 \rightarrow E(\mathbb{F}_q)$ is admissible.*

Remark 1 is still valid when h_{sSWU} is replaced by any encoding implementable with the cost of extracting one square root in \mathbb{F}_q . We chose h_{sSWU} , because it is the most universal among such encodings known in the literature. In particular, this encoding is relevant even if N is a prime (that is the cofactor equals 1), which is the case for many classical elliptic curves. Note that for $q \equiv 11 \pmod{12}$ curves of j -invariants 0, 1728 are supersingular in compliance with [14, §24.2.1.c]. Since such curves pose special challenges for security by virtue of [4, Remark 2.22], the map h_{sSWU} does not have restrictions in the current context.

There is a lot of standardized elliptic curves over fields \mathbb{F}_q such that $q \equiv 11 \pmod{12}$. It is readily checked that this condition is fulfilled, e.g., for the French curve FRP256v1 [25], for the curves P-192, P-384, and Curve448-Goldilocks from NIST SP 800-186 [26, §4.2.1] as well as for all Russian curves [27, Appendix B] except for id-GostR3410-2001-CryptoPro-B-ParamSet. Possibly, Remark 1 can be generalized to the case $q \equiv 2 \pmod{3}$, $q \equiv 5 \pmod{8}$ when a square root is still expressed via one exponentiation (see, e.g., [5, Appendix I.2]). However we did not find standardized curves over such fields, hence we decided to stop in order not to complicate the text.

References

- [1] Koshelev D., *Indifferentiable hashing to ordinary elliptic \mathbb{F}_q -curves of $j = 0$ with the cost of one exponentiation in \mathbb{F}_q* , <https://eprint.iacr.org/2021/301>, accepted in Designs, Codes and Cryptography, 2021.
- [2] Koshelev D., *Faster indifferentiable hashing to elliptic \mathbb{F}_{q^2} -curves*, <https://eprint.iacr.org/2021/678>, 2021.

- [3] Koshelev D., *Optimal encodings to elliptic curves of j -invariants 0*, 1728, <https://eprint.iacr.org/2021/1034>, 2021.
- [4] El Mrabet N., Joye M., *Guide to Pairing-Based Cryptography*, Cryptography and Network Security Series, Chapman and Hall/CRC, New York, 2017.
- [5] Faz-Hernandez A. et al., *Hashing to elliptic curves*, <https://datatracker.ietf.org/doc/draft-irtf-cfrg-hash-to-curve>, 2021.
- [6] Sakemi Y., Kobayashi T., Saito T., Wahby R. S., *Pairing-friendly curves*, <https://datatracker.ietf.org/doc/draft-irtf-cfrg-pairing-friendly-curves>, 2021.
- [7] Budroni A., Pintore F., “Efficient hash maps to \mathbb{G}_2 on BLS curves”, *Applicable Algebra in Engineering, Communication and Computing*, 2020, 1–21.
- [8] Fuentes-Castaneda L., Knapp E., Rodríguez-Henríquez F., “Faster hashing to \mathbb{G}_2 ”, Selected Areas in Cryptography. SAC 2011, LNCS, **7118**, eds. Miri A., Vaudenay S., Springer, Berlin, Heidelberg, 2012, 412–430.
- [9] Scott M., *Authenticated ID-based key exchange and remote log-in with simple token and PIN number*, <https://eprint.iacr.org/2002/164>, 2002.
- [10] Pereira G., Doliskani J., Jao D., “ x -only point addition formula and faster compressed SIKE”, *Journal of Cryptographic Engineering*, **11**:1 (2021), 57–69.
- [11] Boneh D., Gorbunov S. et al., *BLS signatures*, <https://datatracker.ietf.org/doc/draft-irtf-cfrg-bls-signature>, 2020.
- [12] Brier E. et al., “Efficient indifferentiable hashing into ordinary elliptic curves”, *Advances in Cryptology — CRYPTO 2010*, LNCS, **6223**, eds. Rabin T., Springer, Berlin, Heidelberg, 2010, 237–254.
- [13] Wahby R. S., Boneh D., “Fast and simple constant-time hashing to the BLS12-381 elliptic curve”, *IACR Transactions on Cryptographic Hardware and Embedded Systems*, **2019**:4 (2019), 154–179.
- [14] Cohen H. et al., *Handbook of Elliptic and Hyperelliptic Curve Cryptography*, Discrete Mathematics and Its Applications, **34**, Chapman and Hall/CRC, New York, 2005.
- [15] Supranational, *blst/src/sqrt-addchain.h*, <https://github.com/supranational/blst/blob/c76b5ac69a0044432d16cfd2cce60c93c8b01872/src/sqrt-addchain.h>, 2020.
- [16] Tibouchi M., Kim T., “Improved elliptic curve hashing and point representation”, *Designs, Codes and Cryptography*, **82**:1–2 (2017), 161–177.
- [17] Hao F., *Prudent practices in security standardization*, <https://eprint.iacr.org/2021/839>, 2021.
- [18] Crypto Forum Research Group (CFRG), *PAKE selection process*, <https://github.com/cfrg/pake-selection>, 2020.
- [19] Abdalla M., Haase B., Hesse J., *CPace, a balanced composable PAKE*, <https://datatracker.ietf.org/doc/draft-irtf-cfrg-pace/02>, 2021.
- [20] Krawczyk H., Bourdrez D., Lewi K., Wood C. A., *The OPAQUE asymmetric PAKE protocol*, <https://www.ietf.org/id/draft-irtf-cfrg-opaque-06.html>, 2021.
- [21] Abe M., Okamoto T., “Provably secure partially blind signatures”, *Advances in Cryptology — CRYPTO 2000*, LNCS, **1880**, eds. Bellare M., Springer, Berlin, Heidelberg, 2000, 271–286.
- [22] Icart T., “How to hash into elliptic curves”, *Advances in Cryptology — CRYPTO 2009*, LNCS, **5677**, eds. Halevi S., Springer, Berlin, Heidelberg, 2009, 303–316.
- [23] Farashahi R. R. et al., “Indifferentiable deterministic hashing to elliptic and hyperelliptic curves”, *Mathematics of Computation*, **82**:281 (2013), 491–512.
- [24] Koshelev D., *Faster point compression for elliptic curves of j -invariant 0*, <https://eprint.iacr.org/2020/010>, accepted in *Mathematical Aspects of Cryptography*, 2020.
- [25] Agence Nationale de la Sécurité des Systèmes d’Information (ANSSI), *Avis relatif aux paramètres de courbes elliptiques définis par l’Etat français*, <https://www.legifrance.gouv.fr/jorf/id/JORFTEXT000024668816>, 2011.

- [26] Chen L., Moody D., Regenscheid A., Randall K., *Recommendations for discrete logarithm-based cryptography: Elliptic curve domain parameters (Draft NIST Special Publication 800-186)*, <https://csrc.nist.gov/publications/detail/sp/800-186/draft>, 2019.
- [27] Alekseev E. K., Nikolaev V. D., Smyshlyaev S. V., “On the security properties of Russian standardized elliptic curves”, *Mathematical Aspects of Cryptography*, **9:3** (2018), 5–32.